

A K -HOMOLOGICAL APPROACH TO THE QUANTIZATION COMMUTES WITH REDUCTION PROBLEM

YANLI SONG

ABSTRACT. In [15], Kasparov defined a distinguished K -homology fundamental class, so called the Dirac element. We prove a localization formula for the Dirac element in K -homology of crossed product of C^* -algebras. Then we define the quantization of Hamiltonian G -spaces as a push-forward of the Dirac element. With this, we develop a K -homological approach to the quantization commutes with reduction theorem.

CONTENTS

1.	Introduction	1
2.	Analytic K -Homology of C^* -algebras	4
3.	Localization of Dirac Element	9
4.	Dirac Element of Vector Bundles	13
5.	The Cubic Dirac Operator	15
6.	Quantization of Hamiltonian G -Spaces	19
	References	21

1. INTRODUCTION

The purpose of this paper is to study the quantization commutes with reduction principle, which was formulated by Guillemin and Sternberg [8]. There are a number of approaches to the solution such as the geometric proof in [25] and analytic proofs in [33, 28]. By making compromises to both geometry and analysis, we express the quantization commutes with reduction problem in the framework of K -homology. Besides providing a simplified proof, the techniques we used can be adapted to more general situations such as the quasi-Hamiltonian spaces [1], which is our original motivation.

Let G be a compact Lie group, and M a compact G -manifold with a G -equivariant Spin^c -structure. We denote by S_M its spinor bundle. Regarded as a Hilbert bi-module, the bundle S_M induces a push-forward map in K -homology

$$\phi_{S_M} : KK_0^G(C_\tau(M), \mathbb{C}) \rightarrow KK_0^G(\mathbb{C}, \mathbb{C}), \quad (1.1)$$

where $C_\tau(M)$ is the C^* -algebra of continuous sections of the complex Clifford bundle $\text{Cliff}(TM)$. The K -homology on the right-hand side is isomorphic to the representation ring $R(G)$ and the one on the left-hand side contains a distinguished element:

$$[d_M] \in KK_0^G(C_\tau(M), \mathbb{C}),$$

the *Dirac element* [15], defined using the de Rham differential operator d . The push-forward index of the Spin^c -manifold M is defined by:

$$\phi_{S_M}([d_M]) \in KK_0^G(\mathbb{C}, \mathbb{C}) \cong R(G).$$

It coincides with the G -index of the associated Spin^c -Dirac operator on M .

We are particularly interested in the case when G is a compact connected Lie group and (M, ω) is a compact symplectic G -manifold. We also assume that (M, ω) is *pre-quantizable*, that is, there exists a G -equivariant Hermitian line bundle L with a Hermitian connection ∇^L such that

$$\frac{\sqrt{-1}}{2\pi}(\nabla^L)^2 = \omega.$$

Let S_M be the spinor bundle obtained by coupling the pre-quantum line bundle L with the spinor bundle determined by ω . We can formulate the *quantization* of the Hamiltonian G -space (M, ω) as the image of the Dirac element $[d_M]$ under the push-forward map ϕ_{S_M} [26]. We denote it by $Q(M, \omega) \in R(G)$.

When the action of G on M is Hamiltonian, there is a *moment map*

$$\mu : M \rightarrow \mathfrak{g}^*,$$

where $\mathfrak{g}^*$ is the dual of the Lie algebra $\mathfrak{g}$ of G . If 0 is a regular value of μ , then G acts on $\mu^{-1}(0)$ locally freely. For simplicity, we assume that the action is free. We can define the symplectic reduction [36]:

$$M_0 = \mu^{-1}(0)/G, \quad L_0 = (L|_{\mu^{-1}(0)})/G,$$

where M_0 is a symplectic manifold with a symplectic 2-form ω_0 induced from ω . As before, the quantization of the reduced space (M_0, ω_0) is defined as the push-forward of the Dirac element $[d_{M_0}]$ under $\phi_{S_{M_0}}$, denoted by $Q(M_0, \omega_0) \in \mathbb{Z}$.

The quantization commutes with reduction theorem states as follow:

Theorem 1.1. *Let G be a compact connected Lie group, and (M, ω) a compact pre-quantized Hamiltonian G -manifold, with moment map $\mu : M \rightarrow \mathfrak{g}^*$. If 0 is a regular value of μ , then*

$$Q(M, \omega)^G = Q(M_0, \omega_0).$$

Remark 1.2. Guillemin and Sternberg [8] had proved a quantization commutes with reduction theorem for Kähler manifolds. They conjectured the above result for Dirac operators on general symplectic manifolds, see [6]. For the case that G is abelian, the conjecture was solved by Vergne[34] and Meinrenken [24] independently. The general case was first proved by Meinrenken [25] using symplectic surgery, and re-proved by Tian-Zhang [33] using analytic techniques and by Paradan [28] using transversally elliptic operators. After these

results, the quantization commutes with reduction theorem has been generalized in various directions. For example, Ma-Zhang [21, 22] solved an extended version of Vergne’s conjecture [35] on compact Lie group acting on non-compact manifolds. Later on, Paradan [29] gave a different proof. For the case when both manifolds and Lie groups are non-compact, one can find related works by Hochs, Landsman, Mathai and Zhang [12, 11, 19, 23]. At last Paradan and Vergne [30] proved a quantization commutes with reduction formula for Spin^c -manifold very recently.

We now turn to an introduction of our method. We first extend the push-forward map (1.1) to the K -homology of crossed products of C^* -algebras:

$$\hat{\phi}_{S_M} : KK_0(C^*(G, C_\tau(M)), \mathbb{C}) \rightarrow KK_0(C^*(G), \mathbb{C}). \quad (1.2)$$

The target of this map can be identified with $\hat{R}(G) = \text{Hom}_{\mathbb{Z}}(R(G), \mathbb{Z})$, that is, the group of possibly infinite $\mathbb{Z}$ -linear combination $\sum a_\gamma \cdot V_\gamma$ of irreducible representations.

Following an idea of Witten [38], we perturb the de Rham differential operator d by a G -invariant vector field X and define a perturbed Dirac element. We show that the corresponding perturbed Dirac element localizes to a G -invariant open neighborhood U of the vanishing set $\{X = 0\} \subseteq M$.

In the case of Hamiltonian G -space (M, ω) , our method is a variation of Paradan’s approach [28]. We take X to be the Hamiltonian vector field of the norm square of the moment map and U a G -invariant small open neighborhood of $\{X = 0\}$. The submanifold U is disconnected in general. By the symplectic cross-section theorem, every connected component U_α has the geometric structure:

$$U_\alpha \cong G \times_{H_\alpha} V_\alpha, \quad (1.3)$$

where $H_\alpha \subseteq G$ is a sub-group of G of equal rank, and V_α is Hamiltonian H_α -manifold. The tangent bundle TU_α splits equivariantly into “vertical direction” and “horizontal direction”. Hence, so is the Spin^c -structure on U_α and its induced push-forward map.

To prove the quantization commutes with reduction theorem, we need to show that the push-forward index of a connected component U_α doesn’t contribute to the trivial G -representation unless it contains $\mu^{-1}(0)$. By the splitting (1.3) and the Dirac induction formula given by Kostant [18], it is enough to show that the push-forward index of the smaller manifold V_α has zero multiplicity at the trivial H_α -representation.

We then repeat the localization to the Hamiltonian H_α -manifold V_α and reduce the problem to the case when the group is abelian and the manifold is the total space of some vector bundle by induction. With some computation in Section 4, we show that the proof of the quantization commutes with reduction theorem can be localized to a small neighborhood of $\mu^{-1}(0)$.

Acknowledgments: The author benefited a lot from discussions from Victor Guillemin, Nigel Higson and Eckhard Meinrenken. The author also would like to thank Michele Vergne

and Paul-Emile Paradan for reading the first version of this paper and giving some useful advice.

2. ANALYTIC K-HOMOLOGY OF C^* -ALGEBRAS

In this section we give a brief review on Kasparov's approach to analytic K -homology [14, 15].

Definition 2.1. Let A be a separable $\mathbb{Z}_2$ -graded C^* -algebra. A *Fredholm module* over A is a pair $(\mathcal{H}, \mathcal{F})$:

- a separable $\mathbb{Z}_2$ -graded Hilbert space $\mathcal{H}$ with an even $*$ -homomorphism:

$$\rho : A \rightarrow \mathbb{B}(\mathcal{H});$$

- an odd operator $\mathcal{F} \in \mathbb{B}(\mathcal{H})$ such that for all $a \in A$

$$(\mathcal{F} - \mathcal{F}^*)\rho(a) \in \mathbb{K}(\mathcal{H}), (\mathcal{F}^2 - 1)\rho(a) \in \mathbb{K}(\mathcal{H}), [\rho(a), \mathcal{F}] \in \mathbb{K}(\mathcal{H}).$$

Here $\mathbb{K}(\mathcal{H})$ and $\mathbb{B}(\mathcal{H})$ denote the compact operators and the bounded operators on $\mathcal{H}$ respectively.

Definition 2.2. An *operator homotopy* between two Fredholm modules over A : $(\mathcal{H}, \mathcal{F})$ and $(\mathcal{H}, \mathcal{F}')$, is a norm continuous path of Fredholm modules over A

$$t \mapsto (\mathcal{H}, \mathcal{F}_t), \quad t \in [0, 1],$$

with $\mathcal{F}_0 = \mathcal{F}, \mathcal{F}_1 = \mathcal{F}'$.

Definition 2.3. A *unitary equivalence* between two Fredholm modules over A : $(\mathcal{H}, \mathcal{F})$ and $(\mathcal{H}', \mathcal{F}')$, is given by an even unitary isomorphism $T : \mathcal{H} \rightarrow \mathcal{H}'$, which intertwines the actions of A and the operators $\mathcal{F}, \mathcal{F}'$.

Definition 2.4. The *K -homology* $KK_0(A, \mathbb{C})$ is defined as the quotient of the set of Fredholm modules over A by the equivalence relation generated by operator homotopies and unitary equivalences. The addition is given by direct sum and the *zero module* has zero Hilbert space, zero representation, and zero operator.

We can also define $KK_1(A, \mathbb{C}) = KK_0(C_0(\mathbb{R}) \otimes A, \mathbb{C})$ to make $KK_\bullet(A, \mathbb{C})$ into a $\mathbb{Z}_2$ -graded homology theory of C^* -algebras. In a special case when $A = \mathbb{C}$, we have that

$$KK_0(\mathbb{C}, \mathbb{C}) \cong \mathbb{Z}, \quad KK_1(\mathbb{C}, \mathbb{C}) = 0.$$

The following two examples are important in this paper.

Example 2.5. Suppose that (M, g) is an even dimensional complete Riemannian manifold. Let $\text{Cliff}(TM)$ be the bundle of complex Clifford algebras over M with standard $\mathbb{Z}_2$ -grading, and $C_\tau(M)$ the $\mathbb{Z}_2$ -graded C^* -algebra consisting of all the continuous sections of $\text{Cliff}(TM)$ vanishing at infinity. For any $e \in TM$, let $e^* \in T^*M$ corresponds to e via g . The action of e on the exterior algebra of cotangent bundle $\Lambda^*(T^*M)$ is given by the formula:

$$c(e) := \epsilon(e^*) - \iota(e), \quad (2.1)$$

where ϵ is the exterior multiplication and ι is the contraction. This induces an action of $C_\tau(M)$ on $\Lambda^*(T^*M)$.

Let d be the de Rham differential, d^* its dual, and $\mathcal{D} = d + d^*$. The completeness of M ensures that $\mathcal{D}$ is essentially self-adjoint so that we can apply functional calculus to $\mathcal{D}$. The pair

$$\mathcal{H} = \Gamma_{L^2}(M, \Lambda^*(T^*M)), \quad \mathcal{F} = \frac{\mathcal{D}}{\sqrt{\mathcal{D}^2 + 1}} \quad (2.2)$$

gives a Fredholm module over $C_\tau(M)$ and determines an element

$$[d_M] \in KK_0(C_\tau(M), \mathbb{C}).$$

We call this element the *Dirac element* after Kasparov [15, Lemma 4.2].

If M is odd dimensional, one can similarly construct the Dirac element which lives in $KK_1(C_\tau(M), \mathbb{C})$.

Definition 2.6. Let M be a manifold and V an even-dimensional Euclidean vector bundle over M . A *Spin^c-structure* for V is a $\mathbb{Z}_2$ -graded Hermitian vector bundle S over M together with an even isomorphism of $*$ -algebra bundle:

$$c : \text{Cliff}(V) \xrightarrow{\cong} \text{End}(S).$$

In particular, if $V = TM$ has a *Spin^c-structure*, then we say M is a *Spin^c-manifold* and S is its associated *spinor bundle*.

Example 2.7. Let M be an even-dimensional Riemannian manifold, equipped with a *Spin^c-structure*. We denote by S_M its spinor bundle. By choosing a connection ∇ on S_M , we can define a *Spin^c-Dirac operator*

$$D_M = \sum_{i=1}^{\dim(M)} c(e_i) \nabla_{e_i}, \quad (2.3)$$

where $\{e_i\}$ is an orthonormal basis for TM . If M is complete, then we can choose ∇ in a way such that the operator D_M is essentially self-adjoint. By functional calculus, we form

$$\mathcal{H} = \Gamma_{L^2}(M, \Lambda^*(T^*M)), \quad \mathcal{F} = \frac{D_M}{\sqrt{D_M^2 + 1}}. \quad (2.4)$$

Let $C_0(M)$ be the C^* -algebra of continuous functions on M vanishing at infinity, which acts on $\mathcal{H}$ by pointwise multiplication. The pair $(\mathcal{H}, \mathcal{F})$ in (2.4) gives a Fredholm module over $C_0(M)$ and determines an element $[D_M] \in KK_0(C_0(M), \mathbb{C})$.

In the case when M is compact, the map $p : M \rightarrow pt$ collapsing M to a point induces a push-forward map:

$$p_* : KK_\bullet(C(M), \mathbb{C}) \rightarrow KK_\bullet(\mathbb{C}, \mathbb{C}).$$

The integer

$$p_*([D_M]) \in KK_\bullet(\mathbb{C}, \mathbb{C}) \cong \mathbb{Z}$$

equals the index of the *Spin^c-Dirac operator* D_M [10, Chapter 11].

Remark 2.8. Suppose that (M, g) is a compact manifold with a regular boundary in the sense that there is a metric preserving diffeomorphism ϕ between a neighborhood V of ∂M in M and $\partial M \times [0, 1)$. Then $U = \text{Int}(M)$ is a non-complete manifold. However, we can extend the definition of Dirac element to this case. Let $\gamma : U \rightarrow \mathbb{R}$ be a smooth function such that for

$$\gamma(\phi^{-1}(m, t)) = t^2, \quad m \in \partial M, \quad t \leq \frac{1}{2},$$

and

$$\gamma(\phi^{-1}(m, t)) > \frac{1}{4}, \quad m \in \partial M, \quad t > \frac{1}{2},$$

and $\gamma \equiv 1$ on $U \setminus V$. With the rescaled metric $\tilde{g} := \frac{1}{\gamma^2} \cdot g$, $(U, \tilde{g})$ is a complete Riemannian manifold. In addition, the new Clifford action is given by $\tilde{c} = \frac{1}{\gamma} \cdot c$. As the construction in Example 2.5, we can define the Dirac element

$$[d_U] \in KK_{\bullet}(C_{\tau}(U), \mathbb{C}).$$

The K -homology groups $KK_{\bullet}(A, \mathbb{C})$ are functorial not only with respect to C^* -algebra homomorphisms, but more generally with respect to *Morita morphisms* which we shall now describe.

Definition 2.9. If B is a $\mathbb{Z}_2$ -graded C^* -algebra, then a (right) *pre-Hilbert B -module* is a $\mathbb{Z}_2$ -graded complex Banach space $\mathcal{E}$ equipped with a (right) B -action and a B -valued inner product $\langle \cdot, \cdot \rangle_B : \mathcal{E} \times \mathcal{E} \rightarrow B$, which is compatible with the $\mathbb{Z}_2$ -grading, and linear in the second variable and conjugate-linear in the first variable. Moreover, the inner product satisfies

- $(\langle \xi, \eta \rangle_B)^* = \langle \eta, \xi \rangle_B$;
- $\langle \eta, \xi \rangle_B b = \langle \eta, \xi \cdot b \rangle_B$;
- $\|\xi\|^2 = \|\langle \xi, \xi \rangle_B\|$

for all $\xi, \eta \in \mathcal{E}, b \in B$. The norm completion of $\mathcal{E}$ is a *Hilbert B -module*.

The Hilbert $\mathbb{C}$ -modules are precisely Hilbert spaces.

Definition 2.10. Let $\mathcal{E}$ be a Hilbert B -module. We say that a linear map $T : \mathcal{E} \rightarrow \mathcal{E}$ is *adjointable* if there exists a map $T^* : \mathcal{E} \rightarrow \mathcal{E}$ such that

$$\langle T\xi, \eta \rangle_B = \langle \xi, T^*\eta \rangle_B.$$

The space of adjointable maps is denoted by $\mathcal{L}_B(\mathcal{E})$.

Let A, B be two $\mathbb{Z}_2$ -graded C^* -algebras.

Definition 2.11 ([32]). A *Hilbert A - B bi-module* is a pair $(\mathcal{E}, \psi)$ in which $\mathcal{E}$ is a Hilbert B -module and $\psi : A \rightarrow \mathcal{L}_B(\mathcal{E})$ is a $*$ -representation of A on $\mathcal{E}$. Two Hilbert A - B bi-modules $(\mathcal{E}_i, \psi_i), i = 1, 2$ are called *unitarily equivalent* if there is an isomorphism $U : \mathcal{E}_1 \rightarrow \mathcal{E}_2$ preserving the B -valued inner product and $\mathbb{Z}_2$ -grading, and commuting with ψ . In addition, we say that $(\mathcal{E}, \psi)$ is *non-degenerate* if $\psi(A)\mathcal{E} = \mathcal{E}$. *Morita morphisms* between A and B are given by non-degenerate Hilbert A - B bi-modules.

Suppose that $(\mathcal{E}, \psi)$ is a Hilbert A - B bi-module. For any element $[(\mathcal{H}, \mathcal{F})] \in KK_\bullet(B, \mathbb{C})$, where $\mathcal{H}$ is a Hilbert space with left B -module structure. Let $\mathcal{E} \otimes_B \mathcal{H}$ be the norm completion of the algebraic tensor product of $\mathcal{E}$ and $\mathcal{H}$ over B , which is a Hilbert space with a $*$ -representation of A . It is more elaborate to construct the bounded operator $\hat{\mathcal{F}}$ on $\mathcal{E} \otimes_B \mathcal{H}$. We refer to [14, 10] for the construction. Hence, we define a map

$$\phi_{\mathcal{E},*} : KK_\bullet(B, \mathbb{C}) \rightarrow KK_\bullet(A, \mathbb{C}) : [(\mathcal{H}, \mathcal{F})] \mapsto [(\mathcal{E} \otimes_B \mathcal{H}, \hat{\mathcal{F}})]. \quad (2.5)$$

In the viewpoint of Kasparov bivariant K -theory, $[(\mathcal{E} \otimes_B \mathcal{H}, \hat{\mathcal{F}})]$ is the *Kasparov product* of $(\mathcal{H}, \mathcal{F})$ and $(\mathcal{E}, 0)$.

Theorem 2.12. [14] *K -homology is functorial with respect to Morita morphisms.*

Let M be an even-dimensional Riemannian Spin^c -manifold with spinor bundle S_M . For any $v, w \in \Gamma(S_M)$, the formula

$$\langle v, w \rangle := w \cdot v^* \in S_M \otimes S_M^* \cong \text{End}(S_M),$$

defines a $\text{End}(S_M)$ -valued inner product. Moreover, $C_0(M)$ acts on $\Gamma(\text{End}(S_M))$ by point-wise multiplication. By Definition 2.6, we have that

$$\Gamma(\text{End}(S_M)) \cong C_\tau(M),$$

preserving the $\mathbb{Z}_2$ -grading and involution. Therefore, the spinor bundle S_M may be viewed as a Hilbert $C_0(M)$ - $C_\tau(M)$ bi-module.

Theorem 2.13 ([14, 15]). *If M is a complete Riemannian Spin^c -manifold with spinor bundle S_M , then the push-forward map*

$$\Phi_{S_M} : KK_\bullet(C_\tau(M), \mathbb{C}) \rightarrow KK_\bullet(C_0(M), \mathbb{C}),$$

maps $[d_M]$ to $[D_M]$ (see Example 2.5 and 2.7).

Proof. One can find a proof in [26, Proposition C.1]. □

Definition 2.14. For a compact Spin^c -manifold M with spinor bundle S_M , we can define the *push-forward index map*:

$$\phi_{S_M} : KK_\bullet(C_\tau(M), \mathbb{C}) \rightarrow KK_\bullet(\mathbb{C}, \mathbb{C}) \quad (2.6)$$

as a composition:

$$KK_\bullet(C_\tau(M), \mathbb{C}) \xrightarrow{\Phi_{S_M}} KK_\bullet(C(M), \mathbb{C}) \xrightarrow{p_*} KK_\bullet(\mathbb{C}, \mathbb{C}).$$

Remark 2.15. Any two spinor bundles $S_M, \tilde{S}_M$ for M are related by a $\mathbb{Z}_2$ -graded line bundle

$$L = \text{Hom}_{\text{Cliff}(TM)}(S_M, \tilde{S}_M).$$

The two maps $\phi_{S_M}, \phi_{\tilde{S}_M}$ may be different unless the line bundle L can be trivialized.

Let G be a compact Lie group. Everything that we have just discussed generalizes to the G -equivariant case.

Definition 2.16. A $\mathbb{Z}_2$ -graded G - C^* -algebra A is a $\mathbb{Z}_2$ -graded C^* -algebra equipped with a $*$ -homomorphism

$$\alpha : G \rightarrow \text{Aut}(A) : s \mapsto \alpha_s,$$

such that $s \mapsto \alpha_s$ is strongly continuous and preserving the grading.

For any $\mathbb{Z}_2$ -graded G - C^* -algebra A , we can define the G -equivariant K -homology $KK_\bullet^G(A, \mathbb{C})$ with straightforward modification. In particular, if $A = \mathbb{C}$ with trivial G -action, we have that

$$KK_0^G(\mathbb{C}, \mathbb{C}) \cong R(G), \quad KK_1^G(\mathbb{C}, \mathbb{C}) = 0.$$

For more details of equivariant KK -theory, we refer to [15, Section 2].

There is a descent operation which transfers the equivariant KK -theory to the non-equivariant theory. It involves the notion of crossed products of C^* -algebras. We will give a rapid review of the basic construction [37].

If A is a $\mathbb{Z}_2$ -graded G - C^* -algebra, then the space $C(G, A)$ of continuous A -valued functions on G becomes a $\mathbb{Z}_2$ -graded $*$ -algebra with respect to convolution and involution defined by

$$(f * g)(s) = \int_G f(t) \alpha_t(g(t^{-1}s)) dt,$$

and

$$f^*(s) = \alpha_s(f(s^{-1}))^*.$$

A *covariant representation* of A is an even $*$ -representation of A on a $\mathbb{Z}_2$ -graded Hilbert space, together with a compatible unitary representation of G on the same space, that is

$$\pi : A \rightarrow \mathcal{L}(\mathcal{H}), \quad \pi : G \rightarrow U(\mathcal{H}).$$

It determines a $*$ -representation of $C(G, A)$ by the formula:

$$\langle \pi(f)v, w \rangle = \int_G \langle \pi(f(s))\pi(s)v, w \rangle ds, \quad v, w \in \mathcal{H}.$$

Definition 2.17. We define the *crossed product* $C^*(G, A)$ as the completion of $C(G, A)$ under the norm:

$$\|f\| = \sup\{\|\pi(f)\| : \pi \text{ is a covariant representation of } A\}.$$

If $A = \mathbb{C}$, then we get the group C^* -algebra $C^*(G)$. There is a Morita equivalence:

$$C^*(G) \cong \bigoplus_{\gamma \in \text{Irrep}(G)} \text{End}(V_\gamma).$$

It follows that

$$KK_0(C^*(G), \mathbb{C}) \cong \hat{R}(G), \quad KK_1(C^*(G), \mathbb{C}) = 0.$$

We can extend the construction of crossed product to Morita morphisms between C^* -algebras. Let A, B be two $\mathbb{Z}_2$ -graded G - C^* -algebra and $(\mathcal{E}, \psi)$ a Hilbert A - B bi-module, equipped with a strongly continuous homomorphism $u : G \rightarrow \text{Aut}(\mathcal{E})$ such that

$$\langle u_s(\xi), u_s(\eta) \rangle_B = s \cdot (\langle \xi, \eta \rangle_B),$$

and

$$u_s(\xi \cdot b) = u_s(\xi) \cdot (s \cdot b), \quad u_s(\psi(a)\xi) = \psi(s \cdot a) \cdot u_s(\xi),$$

for $\xi, \eta \in \mathcal{E}, a \in A, b \in B, s \in G$. The crossed product $C^*(G, \mathcal{E})$ is a Hilbert $C^*(G, A)$ - $C^*(G, B)$ bi-module, defined as the completion of $C(G, \mathcal{E})$ with respect to the $C^*(G, B)$ -valued inner product:

$$\langle \xi, \eta \rangle(t) = \int_G \langle \xi(s), u_s(\eta(s^{-1}t)) \rangle ds;$$

and with left action of $C(G, A)$ on $C(G, \mathcal{E})$ given by

$$\hat{\psi}(f)(\xi)(t) = \int_G \psi(f(s)) u_s(\xi(s^{-1}t)) ds.$$

The Hilbert bi-module $C^*(G, \mathcal{E})$ gives a Morita morphism between crossed products $C^*(G, A)$ and $C^*(G, B)$, and induces a map:

$$\hat{\phi}_{\mathcal{E},*} : KK_{\bullet}(C^*(G, B), \mathbb{C}) \rightarrow KK_{\bullet}(C^*(G, A), \mathbb{C}). \quad (2.7)$$

Let M be an even-dimensional Riemannian G -manifold. As in Example 2.5, we consider the pair

$$\mathcal{H} = \Gamma_{L^2}(M, \Lambda^*(T^*M)), \quad \mathcal{F} = \frac{\mathcal{D}}{\sqrt{\mathcal{D}^2 + 1}}.$$

For any $f \in C(G, C_{\tau}(M))$, its action on $\mathcal{H}$ is determined by the equation:

$$\langle \pi(f)v, w \rangle = \int_G \langle c(f(s)) \cdot (s \cdot v), w \rangle ds, \quad v, w \in \mathcal{H}, \quad (2.8)$$

where c is the Clifford action. This extends to an action of $C^*(G, C_{\tau}(M))$ on $\mathcal{H}$. One can check that $(\mathcal{H}, \mathcal{F})$ defines a Fredholm module over $C^*(G, C_{\tau}(M))$. When there is no confusion, we still denote by

$$[d_M] \in KK_0(C^*(G, C_{\tau}(M)), \mathbb{C})$$

the element determined by $(\mathcal{H}, \mathcal{F})$. Moreover, if M is compact and carries a G -equivariant Spin^c -structure, then we can extend the push-forward index map:

$$\hat{\phi}_{S_M} : KK_{\bullet}(C^*(G, C_{\tau}(M)), \mathbb{C}) \rightarrow KK_{\bullet}(C^*(G), \mathbb{C}) \cong \hat{R}(G). \quad (2.9)$$

3. LOCALIZATION OF DIRAC ELEMENT

Following an idea of Witten [38], we perturb the de Rham complex by a G -invariant vector field and obtain a localization formula for the Dirac element. In addition, we extend the push-forward index map to non-compact manifolds.

Let G be a compact Lie group and (M, g) a Riemannian G -manifold. Let X be a vector field generated by $\mathfrak{g}$ over M . That is, there exists a G -equivariant map $\psi_X : M \rightarrow \mathfrak{g}$ such that

$$X(m) = \left. \frac{d}{dt} \right|_{t=0} \exp(-t\psi_X(m)) \cdot m, \quad \forall m \in M. \quad (3.1)$$

We assume that the vanishing set $\{X = 0\} \subset M$ is compact. Define a G -invariant function

$$h = \|X\|^2 : M \rightarrow \mathbb{R}$$

and pick a regular value $\epsilon > 0$. Then

$$U = h^{-1}([0, \epsilon)) \subset M$$

is a G -invariant submanifold with a regular boundary. As in Remark 2.8, we can rescale the metric by a G -invariant function $\gamma : U \rightarrow \mathbb{R}$ so that $(U, \tilde{g} = \frac{1}{\gamma^2} \cdot g)$ is a complete Riemannian G -manifold. The Dirac element

$$[d_U] \in KK^G(C^*(G, C_\tau(U)), \mathbb{C}),$$

is represented by

$$\mathcal{H} = \Gamma_{L^2}(U, \wedge^* T^* U), \quad \mathcal{F} = \frac{\mathcal{D}}{\sqrt{\mathcal{D}^2 + 1}}. \quad (3.2)$$

For any $f \in C_\tau(M)$, it acts on $\mathcal{H}$ as the composition:

$$C_\tau(M) \xrightarrow{\text{restriction}} C_\tau(U) \xrightarrow{\frac{1}{\gamma} \cdot c} \mathbb{B}(\mathcal{H}). \quad (3.3)$$

By the formula in (2.8), the above extends to an action of $C^*(G, C_\tau(M))$ on $\mathcal{H}$. However, the pair (3.2) might not be a Fredholm module over $C^*(G, C_\tau(M))$ in general. Instead we introduce the perturbed Dirac element. Let us define a 1-form on U by $X^* := \tilde{g}(X, \cdot)$ and perturb the de Rham differential d and its dual d^* :

$$d_X = d + \epsilon(X^*), \quad d_X^* = d^* + \iota(X). \quad (3.4)$$

Set $\mathcal{D}_X = d_X + d_X^*$.

Lemma 3.1. *One has that*

$$\mathcal{D}_X^2 = (d + d^*)^2 + (\mathcal{L}_X + \mathcal{L}_X^*) + \langle X, X \rangle_{\tilde{g}}.$$

Proof. It follows directly from the Cartan homotopy formula:

$$d\iota_X + \iota_X d = \mathcal{L}_X.$$

□

On the rescaled Riemannian manifold $(U, \tilde{g})$, we define a pair

$$\mathcal{H} = \Gamma_{L^2}(U, \wedge^* T^* U), \quad \mathcal{F}_X = \frac{\mathcal{D}_X}{\sqrt{\mathcal{D}_X^2 + 1}}, \quad (3.5)$$

by replacing $\mathcal{D}$ with $\mathcal{D}_X$.

Lemma 3.2. *If the vanishing set $\{X = 0\}$ is compact, then the pair (3.5) is a Fredholm module over $C^*(G, C_\tau(M))$ and determines an element*

$$[d_{U,X}] \in KK_\bullet(C^*(G, C_\tau(M)), \mathbb{C}).$$

Proof. One has to prove that for any $h \in C^*(G, C_\tau(M))$, the operators

$$h \cdot (1 - \mathcal{F}_X^2), \quad [h, \mathcal{F}_X]$$

are compact on $\mathcal{H}$. We approximate in norm any element $h \in C^*(G, C_\tau(M))$ by finite sums

$$\sum_i e_i \cdot a_i \in C^*(G, C_\tau(M)),$$

where $e_i \in C(G)$ and $a_i \in C_\tau(M)$. Because the operator $\mathcal{F}_X$ is G -invariant, it is enough to prove that the operators

$$a \cdot e \cdot (1 - \mathcal{F}_X^2), \quad e \cdot [a, \mathcal{F}_X] \quad (3.6)$$

are compact.

Let us consider

$$a \cdot (1 + \text{Cas}_{\mathfrak{g}} + \mathcal{D}_X^2)^{-1} = a \cdot (1 + \text{Cas}_{\mathfrak{g}} + \mathcal{D}^2 + \mathcal{L}_X + \mathcal{L}_X^* + \langle X, X \rangle_{\hat{g}})^{-1}, \quad (3.7)$$

where $\text{Cas}_{\mathfrak{g}}$ is the Casimir operator. It is enough to show that the above operator is compact after restricting to every isotypic component $\mathcal{H}^V$ of $\mathcal{H}$, where V is an irreducible G -representation. We notice that both $\text{Cas}_{\mathfrak{g}}$ and $(\mathcal{L}_X + \mathcal{L}_X^*)$ are bounded on $\mathcal{H}^V$, and the function

$$\langle X(m), X(m) \rangle_{\hat{g}} = \frac{1}{\gamma^2} \cdot \langle X(m), X(m) \rangle_g$$

tends to infinity as m tends to the boundary (γ is the rescaling function). By [5, Section 3],

$$a \cdot (1 + \text{Cas}_{\mathfrak{g}} + \mathcal{D}_X^2)^{-1}$$

is a compact operator on $\mathcal{H}^V$ and thus it is compact on $\mathcal{H}$ as well.

On the other hand, we have that

$$\begin{aligned} & a \cdot (1 + \text{Cas}_{\mathfrak{g}} + \mathcal{D}_X^2)^{-1} \cdot e - a \cdot e \cdot (1 + \mathcal{D}_X^2)^{-1} \\ &= a \cdot (1 + \text{Cas}_{\mathfrak{g}} + \mathcal{D}_X^2)^{-1} \cdot ([e, \mathcal{D}_X^2] + \text{Cas}_{\mathfrak{g}} \cdot e) \cdot (1 + \mathcal{D}_X^2)^{-1} \\ &= a \cdot (1 + \text{Cas}_{\mathfrak{g}} + \mathcal{D}_X^2)^{-1} \cdot (\text{Cas}_{\mathfrak{g}} \cdot e) \cdot (1 + \mathcal{D}_X^2)^{-1}. \end{aligned} \quad (3.8)$$

In the last equation, $(1 + \mathcal{D}_X^2)^{-1}$ is automatically bounded and so is $\text{Cas}_{\mathfrak{g}} \cdot e$ ([16, Lemma 6.5]). Therefore, we prove that the operator

$$a \cdot e \cdot (1 + \mathcal{D}_X^2)^{-1} = a \cdot e \cdot (1 - \mathcal{F}_X^2)$$

is compact. One can also show that $e \cdot [a, \mathcal{F}_X]$ is compact by the same argument in [15, Lemma 4.2]. $\square$

For the perturbed Dirac element, we have the following:

Theorem 3.3 (Localization of Dirac element). *Let (M, g) be a complete Riemannian G -manifold. Suppose that X is a vector field generated by $\mathfrak{g}$ over M and $\{X = 0\}$ is compact in M . We have that*

$$[d_M] = [d_{U,X}] \in KK_{\bullet}(C^*(G, C_\tau(M)), \mathbb{C}),$$

where U is a G -invariant open neighborhood of $\{X = 0\}$.

Proof. By definition, the Dirac element $[d_M]$ for (M, g) is represented by

$$\mathcal{H} = \Gamma_{L^2}(M, \wedge^* T^* M), \quad \mathcal{F} = \frac{\mathcal{D}}{\sqrt{\mathcal{D}^2 + 1}}, \quad (3.9)$$

where $\mathcal{D} = d + d^*$. By homotopy, we can also define $[d_M]$ using

$$\mathcal{H} = \Gamma_{L^2}(M, \wedge^* T^* M), \quad \mathcal{F}_X = \frac{\mathcal{D}_X}{\sqrt{\mathcal{D}_X^2 + 1}}, \quad (3.10)$$

where $\mathcal{D}_X = d_X + d_X^*$ is the perturbed differential operator with respect to the metric g .

The perturbed Dirac element $[d_{U,X}]$ for $(U, \tilde{g})$ is given by the pair

$$\tilde{\mathcal{H}} = \Gamma_{L^2}(U, \wedge^* T^* U), \quad \tilde{\mathcal{F}}_X = \frac{\tilde{\mathcal{D}}_X}{\sqrt{\tilde{\mathcal{D}}_X^2 + 1}}, \quad (3.11)$$

where $\tilde{\mathcal{D}}_X = d_X + d_X^*$ is the perturbed differential operator with respect to the rescaled metric $\tilde{g}$. For any irreducible G -representation V , we will show that on the isotypic V -component,

$$[(\mathcal{H}^V, \mathcal{F}_X|_{\mathcal{H}^V})] = [(\tilde{\mathcal{H}}^V, \tilde{\mathcal{F}}_X|_{\tilde{\mathcal{H}}^V})]. \quad (3.12)$$

We choose a relatively compact subset U' such that $\{X = 0\} \subseteq U' \subset U$ and

$$\langle X(m), X(m) \rangle_{\tilde{g}} \gg (\mathcal{L}_X + \mathcal{L}_X^*)|_{\mathcal{H}^V}, \quad m \in U \setminus U'.$$

Without loss of generality, we can also assume that

$$\langle X(m), X(m) \rangle_g \gg (\mathcal{L}_X + \mathcal{L}_X^*)|_{\mathcal{H}^V}, \quad m \in M \setminus U'.$$

In fact, we can always multiply X by some positive function f . Under the above assumptions, the two operators $\mathcal{F}_X|_{\mathcal{H}^V}$ and $\tilde{\mathcal{F}}_X|_{\tilde{\mathcal{H}}^V}$ are invertible outside U' . That is, there exists a positive constant C such that

$$\|\mathcal{F}_X u\| \geq C \cdot \|u\|, \quad \|\tilde{\mathcal{F}}_X v\| \geq C \cdot \|v\|$$

for any $u \in \mathcal{H}^V$ with $\text{Supp } u \subset M \setminus U'$, and $v \in \tilde{\mathcal{H}}^V$ with $\text{Supp } v \subset U \setminus U'$. On the other hand the two operators $\mathcal{F}_X$ and $\tilde{\mathcal{F}}_X$ are homotopic over the relatively compact subset U' . Hence, we proved (3.12). $\square$

Next we generalize the push-forward index to non-compact manifolds. Suppose that M is a possibly non-compact Spin^c G -manifold and X is a G -invariant vector field defined in (3.1). We assume that the vanishing set $\{X = 0\}$ is compact in M . Choose a G -invariant, relatively compact open subset $U \subset M$, containing $\{X = 0\}$. We denote by S_M the G -equivariant spinor bundle for M and S_U the one for U obtained by restriction.

Lemma 3.4. *There exists a G -equivariant embedding $j : U \hookrightarrow N$ into a compact Spin^c G -manifold N , whose G -equivariant spinor bundle S_N can be identified with S_U after restricting to U .*

Proof. Let ρ be a G -invariant function on M with compact support, such that $0 \leq \rho \leq 1$ and $\rho = 1$ on U . We define $\psi : M \times \mathbb{R} \rightarrow \mathbb{R} : (x, t) \mapsto \rho(x) - t^2$. Pick a regular value $0 < c < 1$ of ψ , which always exists by the Sard's theorem. The set $N = \psi^{-1}(c)$ is a compact G -invariant sub-manifold of $U \times \mathbb{R}$. Since M has a G -equivariant Spin^c -structure, so does $M \times \mathbb{R}$ and N . The embedding $j : U \hookrightarrow N$ given by $x \mapsto (x, \sqrt{1-c})$ has the desired properties. $\square$

As in (3.5), we consider the perturbed pair:

$$\mathcal{H} = \Gamma_{L^2}(U, \wedge^* T^* U), \quad \mathcal{F}_X = \frac{\mathcal{D}_X}{\sqrt{\mathcal{D}_X^2 + 1}}. \quad (3.13)$$

Let $C^*(G, C_\tau(N))$ acts on $\mathcal{H}$ factoring through the restriction to U as in (3.3). By Lemma 3.2, the pair $(\mathcal{H}, \mathcal{F}_X)$ defines an element in $KK_\bullet(C^*(G, C_\tau(N)), \mathbb{C})$. Since N is compact,

we define the *push-forward index* of M to be

$$\hat{\phi}_{S_N}([\mathcal{H}, \mathcal{F}_X]) \in KK_{\bullet}(C^*(G), \mathbb{C}) \cong \hat{R}(G). \quad (3.14)$$

Definition 3.5. By the excision property and functoriality of K -homology, the element in (3.14) does not depend on the choices of U , N and the embedding j but only the Spin^c -structure on M and the vector field X . We denote it by

$$\hat{\phi}_{S_M}[d_{M,X}] \in KK_{\bullet}(C^*(G), \mathbb{C}).$$

Remark 3.6. The K -homology index for transversally elliptic operator was introduced by P. Julg [13]. In [16], Kasparov discussed a KK -theoretic approach to transversally elliptic operators in a more general situation. In addition, using the vector field X , Braverman [4] constructed a transversally elliptic operator $D_{M,X}$ on M . The push-forward index

$$\hat{\phi}_{S_M}[d_{M,X}] \in KK_{\bullet}(C^*(G), \mathbb{C}) \cong \hat{R}(G)$$

equals to the G -index of $D_{M,X}$.

4. DIRAC ELEMENT OF VECTOR BUNDLES

The goal of this section is to study the push-forward index of non-compact manifolds which have structures of the total spaces of vector bundles. Throughout this section, we assume that T is a torus. We denote by $\mathfrak{t}$ its Lie algebra and V_{λ} the irreducible representation of T labeled by weight $\lambda \in \mathfrak{t}^*$.

Definition 4.1. Given any element $\chi \in \hat{R}(T)$ and a vector $\alpha \in \mathfrak{t}$, we say that χ is *polarized by α* if the multiplicity of V_{λ} in χ is zero whenever $\langle \lambda, \alpha \rangle < 0$; and we say it is *strictly polarized by α* if the multiplicity of V_{λ} in χ is zero whenever $\langle \lambda, \alpha \rangle \leq 0$.

In particular, if $\chi \in \hat{R}(T)$ is strictly polarized by a vector α , then the multiplicity of the trivial representation in χ must be zero.

Let V be an Euclidean space on which T acts by isometries and preserves only the origin. We fix a vector $\alpha \in \mathfrak{t}$. The Lie derivative $\mathcal{L}_{\alpha}$ is skew-adjoint and the operator

$$J = \frac{\mathcal{L}_{\alpha}}{|\mathcal{L}_{\alpha}|} \in \text{End}(V)$$

defines a T -equivariant almost complex structure on V . The spinor bundle

$$S_V = \wedge^{0,*}(T^*V)$$

defines a T -equivariant Spin^c -structure for V . In addition, we denote by X_{α} the vector field induced by $\alpha \in \mathfrak{t}$.

Lemma 4.2. *The push-forward index*

$$\hat{\phi}_{S_V}([d_{V,X_{\alpha}}]) \in KK_{\bullet}(C^*(T), \mathbb{C}) \cong \hat{R}(T)$$

is polarized by α .

Proof. One can find an explicit formula for the push-forward index in [7, Chapter 7] and [28, Proposition 5.4]. □

More generally suppose that F is an even dimensional Spin^c -manifold and P is a principle bundle over F whose structure group is a compact Lie group H . Let us also assume that H acts on V in such way that it commutes with the T -action and induces a H -action on S_V . In this case, the vector bundle $P \times_H V \rightarrow F$ has a T -equivariant Spin^c -structure, with spinor bundle given by $P \times_H S_V$.

Moreover the total space M of the vector bundle obtains a Spin^c -structure from that of F and V as well. In fact, the tangent bundle TM fits into an exact sequence of vector bundles over M :

$$0 \rightarrow \nu \rightarrow TM \rightarrow \pi^*TF \rightarrow 0,$$

where $\nu = P \times_H TV$ is the vertical tangent bundle, and $\pi : P \rightarrow F$ is the projection map. If we choose a splitting of the sequence then we obtain an isomorphism:

$$TM \cong \nu \oplus \pi^*TF.$$

We define

$$S_M = (P \times_H S_V) \otimes \pi^*S_F, \quad (4.1)$$

and the tangent vector $v \oplus w \in \nu \oplus \pi^*TF$ acts as the operator $v \otimes 1 + 1 \otimes w$ on (4.1). Hence, S_M gives a T -equivariant Spin^c -structure for M . We denote by X_α be the vector field on M induced by $\alpha \in \mathfrak{t}$ as before.

Theorem 4.3. *If the base F is compact, then the push-forward index*

$$\hat{\phi}_{S_M}([d_{M,X_\alpha}]) \in KK_\bullet(C^*(T), \mathbb{C}) \cong \hat{R}(T)$$

is polarized by α .

Proof. One can find an explicit formula for the above push-forward index in [3, Lecture 3]). We now give a simple K -homology approach.

Let us consider a special case in which the principle bundle P is trivial: $P = F \times H$ (in this case we might as well take $H = \{e\}$). Then $M = F \times V$ and the Dirac element $[d_{M,X_\alpha}]$ can be represented by

$$L^2(M, \Lambda^*T^*M) \cong L^2(F, \Lambda^*T^*F) \otimes L^2(V, \Lambda^*T^*V),$$

together operator

$$\mathcal{D}_M = \mathcal{D}_F \otimes 1 + 1 \otimes \mathcal{D}_{V,X_\alpha}$$

where $\mathcal{D}_F = d + d^*$ is the differential operator on F and $\mathcal{D}_{V,X_\alpha} = d_{X_\alpha} + d_{X_\alpha}^*$ is the perturbed differential operator on V . Since $S_M = S_F \boxtimes S_V$, the push-forward map decomposes as well. Hence, we have that

$$\hat{\phi}_{S_M}([d_{M,X_\alpha}]) = \hat{\phi}_{S_F}([d_F]) \times \hat{\phi}_{S_V}([d_{V,X_\alpha}]) \in KK_\bullet(C^*(T), \mathbb{C}).$$

Remembering that T acts trivially on F , one can see that $\hat{\phi}_{S_M}([d_{M,X_\alpha}])$ is polarized by α by Lemma 4.2.

The proof of the general case is similar. We represent the Dirac element $[d_{M, X_\alpha}]$ by the Hilbert space

$$[L^2(P, \Lambda^* \pi^*(T^*F)) \otimes L^2(V, \Lambda^* T^*V)]^H, \quad (4.2)$$

with operator

$$\tilde{D} \otimes 1 + 1 \otimes \mathcal{D}_{X_\alpha},$$

where $\tilde{D}$ is the “lifted” differential operator on P , acting on sections of $\Lambda^* \pi^*(T^*F)$ and acting as identity in the vertical H -direction. This reduces the problem to the first case. $\square$

Remark 4.4. Let $\mathcal{S}$ be an arbitrary T -equivariant spinor bundle for M . It is related with S_M in (4.1) by a T -equivariant $\mathbb{Z}_2$ -graded line bundle

$$L = \text{Hom}_{\text{Cliff}(TM)}(S_M, \mathcal{S}).$$

Recall that T acts trivially on the zero section F . By restricting to F , we have that

$$L|_F = \tilde{L} \otimes \mathbb{C}_\beta,$$

where $\tilde{L}$ is a non-equivariant line bundle on F and $\mathbb{C}_\beta$ is the trivial line bundle on which T acts with weight $\beta \in \mathfrak{t}^*$. If $\langle \beta, \alpha \rangle > 0$, then the push-forward index

$$\hat{\phi}_{\mathcal{S}}([d_{M, X_\alpha}]) \in KK_\bullet(C^*(T), \mathbb{C}) \cong \hat{R}(T)$$

is strictly polarized by α .

When the base F is non-compact, we need additional structures. Let K be another compact Lie group with Lie algebra $\mathfrak{k}$ acting on F . Suppose that F has a K -equivariant Spin^c -structure and P is a K -equivariant principle H -bundle over F . By the above discussion, the product $M = P \times_H V$ is a $K \times T$ -equivariant Spin^c -manifold. Let $\varphi : F \rightarrow \mathfrak{k}$ be a K -equivariant map and define

$$\psi : M \rightarrow \mathfrak{k} \oplus \mathfrak{t} : (x, v) \mapsto (\varphi(x), \alpha). \quad (4.3)$$

The map ψ induces a vector field on M , denoted by X . Under these assumptions, we can use the product model (4.2) to prove the following theorem.

Theorem 4.5. *If the zero set $\{X = 0\}$ is contained in a compact subset of the zero section of M , then the push-forward index:*

$$\hat{\phi}_{S_M}([d_{M, X}]) \in KK_\bullet(C^*(K \times T), \mathbb{C})$$

is polarized by α .

5. THE CUBIC DIRAC OPERATOR

Dirac operators on homogeneous spaces were discussed in a number of references. We will study a specific algebraically defined operator, that is, the cubic Dirac operator after Kostant [18]. The key property of the cubic Dirac operator is that its square equals to the quadratic Casimir operator up to low order terms. We use [27] as our primary reference.

Let G be a compact Lie group and $\mathfrak{g}$ its Lie algebra. We equip $\mathfrak{g}$ with an ad-invariant inner product and fix an orthonormal basis:

$$X_i, \quad i = 1, \dots, \dim(\mathfrak{g}).$$

Let $S_{\mathfrak{g}}$ be a $\mathbb{Z}_2$ -graded irreducible representation of the Clifford algebra $\text{Cliff}(\mathfrak{g})$, together with *Clifford action*:

$$c : \text{Cliff}(\mathfrak{g}) \hookrightarrow \text{End}(S_{\mathfrak{g}}).$$

If $\mathfrak{g}$ is even dimensional, then $\text{Cliff}(\mathfrak{g}) \cong \text{End}(S_{\mathfrak{g}})$. The *spin representation* $\text{ad}^{\mathfrak{g}} : \mathfrak{g} \rightarrow \text{End}(S_{\mathfrak{g}})$ is defined by the formula:

$$\text{ad}^{\mathfrak{g}}(X) := \frac{1}{4} \sum_{i=1}^{\dim(\mathfrak{g})} c([X, X_i])c(X_i), \quad X \in \mathfrak{g}.$$

Definition 5.1. Let V be a G -representation with infinitesimal $\mathfrak{g}$ -action π . For any $q \in \mathbb{R}^+$, we define an operator on $V \otimes S_{\mathfrak{g}}$ by

$$\begin{aligned} D_{\mathfrak{g}}^q &= \sum_{i=1}^{\dim(\mathfrak{g})} (\pi(X_i) \otimes c(X_i) + q \otimes \text{ad}^{\mathfrak{g}}(X_i) \cdot c(X_i)) \\ &= \sum_{i=1}^{\dim(\mathfrak{g})} c(X_i) \cdot (\pi(X_i) + q \cdot \text{ad}^{\mathfrak{g}}(X_i)). \end{aligned} \tag{5.1}$$

Note that the second form resembles a geometric Dirac operator for the connection on the spinor bundle $S_{\mathfrak{g}}$:

$$\nabla_X^q = \pi(X_i) + q \cdot \text{ad}^{\mathfrak{g}}(X_i).$$

Taking $q = \frac{1}{2}$, $\nabla^{\frac{1}{2}}$ is the Levi-Civita connection and $D_{\mathfrak{g}}^{\frac{1}{2}}$ is the usual Riemannian Dirac operator; while $q = \frac{1}{3}$ gives the *cubic Dirac operator*, denoted by $D_{\mathfrak{g}}$.

The cubic Dirac operator $D_{\mathfrak{g}}$ gives an element in the quantum Weil algebra $\mathcal{W}(\mathfrak{g}) := \mathcal{U}(\mathfrak{g}) \otimes \text{Cliff}(\mathfrak{g})$ [2, 27] and has the following interesting property.

Theorem 5.2 ([18]). *The square of the cubic Dirac operator equals*

$$D_{\mathfrak{g}}^2 = 2 \cdot \text{Cas}_{\mathfrak{g}} + \frac{1}{12} \text{tr}(\text{Cas}_{\mathfrak{g}}).$$

Let $P_{G,+} \subset \Lambda^*$ be the set of dominant weights for G . The set of finite dimensional irreducible G -representations is parametrized by $P_{G,+}$. In particular, for any irreducible representation V_{λ} with highest weight $\lambda \in P_{G,+}$, $D_{\mathfrak{g}}^2$ acts on $V_{\lambda} \otimes S_{\mathfrak{g}}$ as the scalar $\|\lambda + \rho_G\|^2$, where ρ_G is the half sum of positive roots of G .

More generally, Kostant introduced the *relative cubic Dirac operator* for pairs of Lie groups. Let $H \subset G$ be a closed subgroup of the equal rank. Using the inner product we write $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{p}$. This decomposition induces isomorphisms:

$$\text{Cliff}(\mathfrak{g}) \cong \text{Cliff}(\mathfrak{h}) \otimes \text{Cliff}(\mathfrak{p}),$$

and

$$S_{\mathfrak{g}} \cong S_{\mathfrak{h}} \otimes S_{\mathfrak{p}}, \tag{5.2}$$

where $S_{\mathfrak{h}}, S_{\mathfrak{p}}$ are $\mathbb{Z}_2$ -graded irreducible modules of $\text{Cliff}(\mathfrak{h})$ and $\text{Cliff}(\mathfrak{p})$. Accordingly the operator $\text{ad}(X)$ breaks up as a sum

$$\text{ad}^{\mathfrak{g}}(X) = \text{ad}^{\mathfrak{h}}(X) + \text{ad}^{\mathfrak{p}}(X), \quad X \in \mathfrak{h}.$$

Let $D'_{\mathfrak{h}}$ be the cubic Dirac operator associated to the $\mathfrak{h}$ -representation on $V \otimes S_{\mathfrak{g}} \cong (V \otimes S_{\mathfrak{p}}) \otimes S_{\mathfrak{h}}$.

Definition 5.3. The *relative cubic Dirac operator* $D_{\mathfrak{g}, \mathfrak{h}}$ is defined as the difference $D_{\mathfrak{g}, \mathfrak{h}} = D_{\mathfrak{g}} - D'_{\mathfrak{h}}$, which can be written out as

$$D_{\mathfrak{g}, \mathfrak{h}} = \sum_{i=1}^{\dim(\mathfrak{p})} (\pi(X_i) \otimes c(X_i) + 1 \otimes \frac{1}{3} \cdot \text{ad}^{\mathfrak{p}}(X_i) \cdot c(X_i)), \quad (5.3)$$

where $\{X_i\}_{i=1}^{\dim(\mathfrak{p})}$ is an orthonormal basis for $\mathfrak{p}$.

For the relative cubic Dirac operator, we have the following theorem.

Theorem 5.4 ([18]). *Let V_{λ} be an irreducible representation of G . On each isotypic H -summand W_{μ} of $V_{\lambda} \otimes S_{\mathfrak{p}}$ labeled by highest weight $\mu \in \Lambda_{H,+}^*$, the square of the relative Dirac operator acts as*

$$D_{\mathfrak{g}, \mathfrak{h}}^2|_{W_{\mu}} = \|\lambda + \rho_G\|^2 - \|\mu + \rho_H\|^2.$$

Suppose that the adjoint H -action on $\mathfrak{p}$ lifts to a H -action on $S_{\mathfrak{p}}$. In this case, the bundle

$$S_{G/H} = G \times_H S_{\mathfrak{p}}^* \rightarrow G/H,$$

where $S_{\mathfrak{p}}^*$ is the dual of the spinor module $S_{\mathfrak{p}}$, gives a G -equivariant Spin structure on the homogeneous space G/H . Given any H -space W , we can form a G -equivariant vector bundle $E_W = G \times_H W$ over G/H . The space of L^2 -sections of the tensor bundle

$$S_{G/H} \otimes E_W = G \times_H (S_{\mathfrak{p}}^* \otimes W) \rightarrow G/H$$

can be identified with

$$\mathcal{H} = [L^2(G) \otimes S_{\mathfrak{p}}^* \otimes W]^H \cong \bigoplus_{\lambda \in P_{G,+}} V_{\lambda} \otimes \text{Hom}_H(V_{\lambda} \otimes S_{\mathfrak{p}}, W),$$

where the isomorphism comes from the Peter-Weyl theorem.

We now define a Dirac operator on $\mathcal{H}$. Since the relative cubic Dirac operator $D_{\mathfrak{g}, \mathfrak{h}}$ is H -equivariant, it restrict to an operator $D_{\mathfrak{g}, \mathfrak{h}}^{V_{\lambda}}$ on

$$\text{Hom}_H(V_{\lambda} \otimes S_{\mathfrak{p}}, W).$$

Tensoring each operator $D_{\mathfrak{g}, \mathfrak{h}}^{V_{\lambda}}$ with the identity operator on V_{λ} , and summing over V_{λ} , one obtains an operator D_W on $\mathcal{H}$.

In addition, the Hilbert space $\mathcal{H}$ carries a $\mathbb{Z}_2$ -grading coming from that on $S_{\mathfrak{p}}$. We write $\mathcal{H} = \mathcal{H}^+ \oplus \mathcal{H}^-$ and the operator D_W anti-commutes with the $\mathbb{Z}_2$ -grading. Since D_W is equivariant, the kernel $\ker(D_W) \in R(G)$.

Definition 5.5. We define the *Dirac induction*

$$\mathcal{I}_H^G : R(H) \rightarrow R(G) \quad (5.4)$$

to be the map

$$W \mapsto \text{Index}(D_W) = \ker(D_W) \cap \mathcal{H}^+ - \ker(D_W) \cap \mathcal{H}^-.$$

We can describe the Dirac induction map in terms of weights.

Theorem 5.6 ([18]). *Let W_λ be an irreducible representation of H with highest weight $\lambda \in P_{H,+}$. If there exists a $w \in W_G$ in the Weyl group (necessarily unique) such that*

$$\mu + \rho_G = w \cdot (\lambda + \rho_H)$$

for some $\mu \in P_{G,+}$, then the Dirac induction takes W_λ to the $(-1)^w \cdot V_\mu$; Otherwise, the Dirac induction takes W_λ to 0.

The Dirac induction in (5.4) can be extended to a map:

$$\hat{\mathcal{I}}_H^G : \hat{R}(H) \rightarrow \hat{R}(G).$$

Remark 5.7. Landweber [20] and Posthuma [31] generalized the construction of relative cubic Dirac operators to the loop group setting, in which they obtained different generalizations of the Weyl-Kac formula.

Let us briefly compare with *holomorphic induction*. Suppose that $\mathcal{O} = G/H$ is an orbit of coadjoint G -action on $\mathfrak{g}^*$ and U is a H -equivariant Spin^c -manifold, with spinor bundle S_U . It is well-known that $\mathcal{O}$ is a complex G -manifold with spinor bundle

$$S_{\mathcal{O}} = G \times_H (S_{\mathfrak{p}}^* \otimes \mathbb{C}_{\rho_G - \rho_H}), \quad (5.5)$$

where $\mathbb{C}_{\rho_G - \rho_H}$ is the complex plane on which H acts with weight $\rho_G - \rho_H$.

The product $M = G \times_H U$ obtains a G -equivariant Spin^c -structure from that of U and $\mathcal{O}$. In fact, the tangent bundle TM splits into vertical direction and horizontal direction:

$$TM \cong (G \times_H TU) \oplus \pi^* T\mathcal{O}.$$

where $\pi : M \rightarrow \mathcal{O}$ is the projection. We form a tensor product

$$S_M = S_\nu \hat{\otimes} \pi^* S_{\mathcal{O}}, \quad (5.6)$$

where $S_\nu = G \times_H S_U$. The tangent vector

$$v \oplus w \in (G \times_H TU) \oplus \pi^* T\mathcal{O}$$

acts as the operator

$$v \hat{\otimes} 1 + 1 \hat{\otimes} w$$

on $S_\nu \hat{\otimes} \pi^* S_{\mathcal{O}}$. Hence, S_M defines a spinor bundle for M .

We define two Dirac-type operators on the Hilbert space

$$\Gamma_{L^2}(M, S_M) \cong \bigoplus_{\lambda \in P_{G,+}} V_\lambda \otimes \text{Hom}_H(V_\lambda \otimes S_{\mathfrak{p}}, \Gamma_{L^2}(U, S_U) \otimes \mathbb{C}_{\rho_G - \rho_H}). \quad (5.7)$$

We first construct a Spin^c -Dirac operator D_M by choosing a connection on the spinor bundle S_M as in Example 2.7.

On the other hand, one can also build a Dirac operator on (5.7) as a sum

$$\tilde{D}_M = D_{\text{ver}} + D_{\text{hor}}.$$

Here the vertical part D_{ver} acts on the factor $\Gamma_{L^2}(U, S_U)$ as the Spin^c -Dirac operator D_U ; and the horizontal part D_{hor} is constructed as in the Dirac induction so that it acts as the relative cubic Dirac operator on the factors $V_\lambda \otimes S_{\mathfrak{p}}$. One can see that $\tilde{D}_M$ is a combination of geometrically defined operator and algebraically defined operator.

The two operators $\tilde{D}_M$ and D_M are identical in the vertical direction; while in the horizontal direction, one is the relative cubic Dirac operator and the other is the Spin^c -Dirac operator. As pointed out in Definition 5.1, these two operators are homotopic. Therefore, the push-forward index $\hat{\phi}_{S_M}([d_M]) \in \hat{R}(G)$ can be obtained by induction from $\hat{\phi}_{S_U}([d_U]) \in \hat{R}(H)$. By Theorem 5.6, we have the following.

Corollary 5.8. *The push-forward index $\hat{\phi}_{S_M}([d_M]) \in \hat{R}(G)$ has the same multiplicity at the trivial representation as $\hat{\phi}_{S_U}([d_U]) \in \hat{R}(H)$.*

6. QUANTIZATION OF HAMILTONIAN G -SPACES

The goal of this section is to re-prove Theorem 1.1. Let us consider a Hamiltonian action by a compact connected Lie group G on a possibly non-compact symplectic manifold (M, ω) . Suppose that (M, ω) is pre-quantizable, with pre-quantum line bundle L and connection ∇^L . We can define the moment map $\mu : M \rightarrow \mathfrak{g}^*$ by the Kostant's formula:

$$\langle \mu, \xi \rangle = \frac{\sqrt{-1}}{2\pi} (\nabla_{X_\xi}^L - \mathcal{L}_\xi), \quad \xi \in \mathfrak{g}$$

where $\mathcal{L}_\xi$ is the Lie derivative and X_ξ is the vector field induced by the infinitesimal action of ξ .

We choose a G -equivariant almost complex structure J so that

$$g(\cdot, \cdot) = \omega(\cdot, J\cdot)$$

defines a Riemannian metric on M . Coupling with the pre-quantum line bundle L , the symplectic 2-form ω determines a spinor bundle

$$S_M = \wedge^{0,*}(T^*M) \otimes L.$$

Fix an Ad-invariant inner product of $\mathfrak{g}$. Let X be the Hamiltonian vector field of the norm square of the moment map $\|\mu\|^2$, which is known as the *Kirwan vector field*. By calculation, we see that

$$X = -J(d\|\mu\|^2)^* = 2 \sum_{i=1}^{\dim(\mathfrak{g})} \mu(\xi_i) \cdot X_{\xi_i}$$

where $\{\xi_i\}$ is an orthonormal basis of $\mathfrak{g}$. From the viewpoint of (3.1), the vector field X is induced by $2 \cdot \mu$. The Kirwan vector field plays an essential role in the work of Tian-Zhang [33] and Paradan [28].

Let U be a G -invariant small open neighborhood of the vanishing set $\{X = 0\}$. In [17], Kirwan showed that $\{X = 0\}$ can be decomposed:

$$\{X = 0\} = \bigsqcup_{\alpha \in \Gamma} G \cdot (M^\alpha \cap \mu^{-1}(\alpha)), \quad (6.1)$$

where Γ is a discrete subset of positive Weyl chamber $\mathfrak{t}_+$. Accordingly U breaks up into different connected components U_α . In particular, U_0 contains $\mu^{-1}(0)$. If $\{X = 0\}$ is compact in M , then we have the localization formula:

$$[d_M] = [d_{U,X}] = \sum_{\alpha \in \Gamma} [d_{U_\alpha, X}] \in KK_0(C^*(G, C_\tau(M)), \mathbb{C}). \quad (6.2)$$

Theorem 1.1 follows directly from the following two theorems.

Theorem 6.1. *If 0 is a regular value of μ , then the multiplicity of the trivial representation in*

$$\hat{\phi}_{S_M|_{U_0}}([d_{U_0, X}]) \in KK_0(C^*(G), \mathbb{C}) \cong \hat{R}(G)$$

equals to $Q(M_0, \omega_0) \in \mathbb{Z}$.

Theorem 6.2. *For each $\alpha \in \Gamma$, if $\alpha \neq 0$, then the element*

$$\hat{\phi}_{S_M|_{U_\alpha}}([d_{U_\alpha, X}]) \in KK_0(C^*(G), \mathbb{C}) \cong \hat{R}(G)$$

doesn't contribute to the multiplicity of the trivial representation.

Remark 6.3. For Theorem 6.1 and 6.2, one can also find different proofs in [28, 33].

We focus on Theorem 6.2 for the rest of this section. Let us recall the symplectic cross-section theorem [9]:

Theorem 6.4. *Let (M, ω) be a symplectic manifold with a moment map $\mu : M \rightarrow \mathfrak{g}^*$ arising from a Hamiltonian G -action. Let α be a point in $\mathfrak{g}^*$ and G_α its isotropy Lie group. There exists a G_α -invariant subset $R_\alpha \subset \mathfrak{g}^*$ containing α such that the cross-section $V_\alpha = \mu^{-1}(R_\alpha)$ is a G_α -invariant symplectic sub-manifold of M , and the set*

$$U_\alpha := G \cdot V_\alpha \cong G \times_{G_\alpha} V_\alpha$$

is diffeomorphic to an open subset in M . Moreover, if M has a pre-quantum line bundle L , then V_α has a pre-quantum line bundle L_α as well, with the property that $L|_{U_\alpha} \cong G \times_{G_\alpha} L_\alpha$. The moment map associated to the G_α -action on V_α is obtained by restriction.

We apply the symplectic cross-section theorem to our case and conclude that each component U_α has the geometric structure:

$$U_\alpha \cong G \times_{G_\alpha} V_\alpha. \quad (6.3)$$

Coupling with the pre-quantum line bundle L_α , the symplectic 2-form on V_α determines a G_α -equivariant Spin^c -structure on V_α with spinor bundle

$$S_{V_\alpha} = \wedge^{0,*}(T^*V_\alpha) \otimes L_\alpha.$$

Together with the complex structure on G/G_α , we obtain a G -equivariant Spin^c -structure on U_α as in (5.6), which is equivalent to that obtained by restricting S_M to U_α . By Corollary 5.8, the push-forward index

$$\hat{\phi}_{S_M|_{U_\alpha}}([d_{U_\alpha, X}]) \in KK_0(C^*(G), \mathbb{C}) \cong \hat{R}(G).$$

has the same multiplicity at the trivial representation as

$$\hat{\phi}_{S_{V_\alpha}}([d_{V_\alpha, X}]) \in KK_0(C^*(G_\alpha), \mathbb{C}) \cong \hat{R}(G_\alpha).$$

Therefore, it is enough to show that $\hat{\phi}_{S_{V_\alpha}}([d_{V_\alpha, X}])$ does not contribute the trivial representation.

The smaller Hamiltonian G_α -manifold V_α has a moment map obtained by restriction. We can apply the localization process again to V_α . By induction, we reduce the problem to the case when $G_\alpha = G$. We can thus assume $G = K \times T_\alpha$, where T_α the torus generated by α . Accordingly the moment map breaks up as a sum:

$$\mu = \mu_{\mathfrak{k}} + \mu_\alpha$$

where $\mu_{\mathfrak{k}}, \mu_\alpha$ are the composition of μ with the projection from $\mathfrak{g}$ to $\mathfrak{k}$ and $\text{Lie}(T_\alpha)$ respectively.

Let F be a K -invariant open neighborhood of $M^\alpha \cap \mu^{-1}(\alpha)$ in M^α and N a G -invariant tubular neighborhood of F in M . In order to apply the Theorem 4.5, we need to deform the moment map μ . We first shrink F and N so that there is a small constant ϵ satisfying:

- (1) $\|\alpha - \mu(n)\| < \epsilon$ for all $n \in N$;
- (2) N^γ is contained in the zero section of N for all $\|\gamma - \alpha\| < \epsilon, \gamma \in \mathfrak{g}$. This is possible because N^α preserves only the zero section.

For $0 \leq t \leq 1$, we define

$$\mu^t(x, v) = \mu_{\mathfrak{k}}(x, t \cdot v) + \mu_\alpha(x, v) : N \rightarrow \mathfrak{g},$$

and X^t the vector field induced by $2\mu^t$. The two conditions above ensures that the vanishing set $\{X^t = 0\}$ is contained in a compact subset of the zero section of N for all $0 \leq t \leq 1$. By homotopy,

$$[d_{N, X}] = [d_{N, X^0}] \in KK_0(C^*(G, C_\tau(M)), \mathbb{C}).$$

By identifying U_α with N , we have the following:

Proposition 6.5. *The element*

$$\hat{\phi}_{S_M|U_\alpha}([d_{U_\alpha, X^0}]) \in KK_0(C^*(G), \mathbb{C}) \cong \hat{R}(G)$$

has zero multiplicity at the trivial representation.

Proof. We can now apply Theorem 4.5 to show that

$$\hat{\phi}_{S_{V_\alpha}}([d_{U_\alpha, X^0}]) \in KK_0(C^*(K \times T_\alpha), \mathbb{C}) \cong \hat{R}(K \times T_\alpha)$$

is polarized by α . Moreover, the restriction of the pre-quantum line bundle L to the T_α -fixed zero section F has a T_α -weight $\alpha \in \mathfrak{k}$. Thus, it provides a “positive factor”. By Remark 4.4, $\hat{\phi}_{S_{V_\alpha}}([d_{U_\alpha, X^0}])$ is strictly polarized by α . $\square$

REFERENCES

- [1] Anton Alekseev, Anton Malkin, and Eckhard Meinrenken. Lie group valued moment maps. *J. Differential Geom.*, 48(3):445–495, 1998.

- [2] Anton. Alekseev and Eckhard. Meinrenken. The non-commutative Weil algebra. *Invent. Math.*, 139(1):135–172, 2000.
- [3] Michael Francis Atiyah. *Elliptic operators and compact groups*. Lecture Notes in Mathematics, Vol. 401. Springer-Verlag, Berlin, 1974.
- [4] Maxim Braverman. Index theorem for equivariant Dirac operators on noncompact manifolds. *K-Theory*, 27(1):61–101, 2002.
- [5] Mikhael Gromov and H. Blaine Lawson, Jr. Positive scalar curvature and the Dirac operator on complete Riemannian manifolds. *Inst. Hautes Études Sci. Publ. Math.*, (58):83–196 (1984), 1983.
- [6] Victor Guillemin. Reduced phase spaces and Riemann-Roch. In *Lie theory and geometry*, volume 123 of *Progr. Math.*, pages 305–334. Birkhäuser Boston, Boston, MA, 1994.
- [7] Victor Guillemin, Viktor Ginzburg, and Yael Karshon. *Moment maps, cobordisms, and Hamiltonian group actions*, volume 98 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2002. Appendix J by Maxim Braverman.
- [8] Victor. Guillemin and Shlomo Sternberg. Geometric quantization and multiplicities of group representations. *Invent. Math.*, 67(3):515–538, 1982.
- [9] Victor Guillemin and Shlomo Sternberg. *Symplectic techniques in physics*. Cambridge University Press, Cambridge, 1984.
- [10] Nigel Higson and John Roe. *Analytic K-homology*. Oxford Mathematical Monographs. Oxford University Press, Oxford, 2000. Oxford Science Publications.
- [11] Peter Hochs. Quantisation commutes with reduction at discrete series representations of semisimple groups. *Adv. Math.*, 222(3):862–919, 2009.
- [12] Peter. Hochs and Klaas. Landsman. The Guillemin-Sternberg conjecture for noncompact groups and spaces. *J. K-Theory*, 1(3):473–533, 2008.
- [13] Pierre Julg. Induction holomorphe pour le produit croisé d’une C^* -algèbre par un groupe de Lie compact. *C. R. Acad. Sci. Paris Sér. I Math.*, 294(5):193–196, 1982.
- [14] Gennadi Kasparov. The operator K -functor and extensions of C^* -algebras. *Izv. Akad. Nauk SSSR Ser. Mat.*, 44(3):571–636, 719, 1980.
- [15] Gennadi Kasparov. Equivariant KK -theory and the Novikov conjecture. *Invent. Math.*, 91(1):147–201, 1988.
- [16] Gennadi Kasparov. K-theoretic index theorems for elliptic and transversally elliptic operators. *Journal of K-theory*, To appear.
- [17] Frances Clare Kirwan. *Cohomology of quotients in symplectic and algebraic geometry*, volume 31 of *Mathematical Notes*. Princeton University Press, Princeton, NJ, 1984.
- [18] Bertram Kostant. A cubic Dirac operator and the emergence of Euler number multiplets of representations for equal rank subgroups. *Duke Math. J.*, 100(3):447–501, 1999.
- [19] N. P. Landsman. Functorial quantization and the Guillemin-Sternberg conjecture. In *Twenty years of Białowieża: a mathematical anthology*, volume 8 of *World Sci. Monogr. Ser. Math.*, pages 23–45. World Sci. Publ., Hackensack, NJ, 2005.
- [20] Gregory D. Landweber. Multiplets of representations and Kostant’s Dirac operator for equal rank loop groups. *Duke Math. J.*, 110(1):121–160, 2001.
- [21] Xiaonan Ma and Weiping Zhang. Geometric quantization for proper moment maps. *C. R. Math. Acad. Sci. Paris*, 347(7-8):389–394, 2009.
- [22] Xiaonan Ma and Weiping Zhang. Geometric quantization for proper moment maps: the Vergne conjecture. *Acta Math.*, 212(1):11–57, 2014.
- [23] Varghese Mathai and Weiping Zhang. Geometric quantization for proper actions. *Adv. Math.*, 225(3):1224–1247, 2010. With an appendix by Ulrich Bunke.

- [24] Eckhard Meinrenken. On Riemann-Roch formulas for multiplicities. *J. Amer. Math. Soc.*, 9(2):373–389, 1996.
- [25] Eckhard Meinrenken. Symplectic surgery and the Spin^c -Dirac operator. *Adv. Math.*, 134(2):240–277, 1998.
- [26] Eckhard. Meinrenken. Twisted K -homology and group-valued moment maps. *Int. Math. Res. Not. IMRN*, (20):4563–4618, 2012.
- [27] Eckhard Meinrenken. *Clifford algebras and Lie theory*, volume 58 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer, Heidelberg, 2013.
- [28] Paul-Émile Paradan. Localization of the Riemann-Roch character. *J. Funct. Anal.*, 187(2):442–509, 2001.
- [29] Paul-Émile Paradan. Formal geometric quantization II. *Pacific J. Math.*, 253(1):169–211, 2011.
- [30] Paul-Émile Paradan and Michèle Vergne. The multiplicities of the equivariant index of twisted Dirac operators. *C. R. Math. Acad. Sci. Paris*, 352(9):673–677, 2014.
- [31] Hessel Posthuma. Dirac induction for loop groups. *Lett. Math. Phys.*, 95(1):89–107, 2011.
- [32] Marc A. Rieffel. Induced representations of C^* -algebras. *Advances in Math.*, 13:176–257, 1974.
- [33] Youliang Tian and Weiping Zhang. An analytic proof of the geometric quantization conjecture of Guillemin-Sternberg. *Invent. Math.*, 132(2):229–259, 1998.
- [34] Michèle Vergne. Quantification géométrique et multiplicités. *C. R. Acad. Sci. Paris Sér. I Math.*, 319(4):327–332, 1994.
- [35] Michèle Vergne. Applications of equivariant cohomology. In *International Congress of Mathematicians. Vol. I*, pages 635–664. Eur. Math. Soc., Zürich, 2007.
- [36] Alan Weinstein. *Lectures on symplectic manifolds*, volume 29 of *CBMS Regional Conference Series in Mathematics*. American Mathematical Society, Providence, R.I., 1979. Corrected reprint.
- [37] Dana P. Williams. *Crossed products of C^* -algebras*, volume 134 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2007.
- [38] Edward Witten. Supersymmetry and Morse theory. *J. Differential Geom.*, 17(4):661–692 (1983), 1982.